

# Indeterministic Quantum Gravity

## II. Refinements and Developments

Vladimir S. MASHKEVICH<sup>1</sup>

*Institute of Physics, National academy of sciences of Ukraine  
252028 Kiev, Ukraine*

### Abstract

This paper is a continuation of the paper [1]. Indeterministic quantum gravity is a theory that unifies general relativity and quantum theory involving indeterministic conception, i.e., quantum jumps. By the same token the theory claims to describe all the universe.

Spacetime is the direct product of cosmic time and space. The state of the universe is given by metric, its derivative with respect to cosmic time, and the number of an energy level. A quantum jump occurs at the tangency of two levels. Equations of motion are the restricted Einstein equation (the cosmic space part thereof) and a probability rule for the quantum jump.

---

<sup>1</sup>E-mail: mash@phys.unit.no

## Introduction

The idea of a possible role of gravity in quantum indeterminism, i.e., in state vector reduction is not new. It has been advanced, among others, by Feynman [2]. A discussion of this role is contained in the paper and the book [3] by Penrose. In the papers [4] there is a calculation of a possible coherence breaking due to metric fluctuations. But it is only in the paper [1] that the idea has been embodied in an indeterministic dynamics. To do this a new physical principle has been advanced — that of cosmic energy determinacy.

This paper, being a continuation of the paper [1], contains some refinements and developments. The most important ones are the following.

Due to quantum jumps, it is only the restricted Einstein equation that holds, namely,  $G^{ij} = (\Psi, T^{ij}\Psi)$ , where  $i, j = 1 = 2 = 3$  correspond to cosmic space.

The state of the universe is  $\omega = (g, g', l)$ , where  $g$  is metric,  $g'$  is its derivative with respect to cosmic time, and  $l$  is the number of an energy level.

A quantum jump occurs at and only at a tangency of two levels.

Equations of motion reduce to the restricted Einstein equation and a probability rule for the quantum jump.

## 1 The universe as a physical system: Spacetime and matter

The universe  $U$  as a physical system is a pair:

$$U = (st, m), \quad (1.1)$$

where  $st$  is spacetime,  $m$  is matter. The spacetime is a triple:

$$st = (M, g, \nabla), \quad (1.2)$$

where  $M$  is a differentiable 4-dimensional manifold,  $g$  is a metric, and  $\nabla$  is the Levi-Civita connection. The matter is a family  $F$  of quantum fields on  $M$ :

$$m = F. \quad (1.3)$$

## 2 The structure of spacetime: Cosmic time and space

The manifold  $M$  is a direct product of two manifolds:

$$M = T \times S, \quad M \ni p = (t, s), \quad t \in T, \quad s \in S. \quad (2.1)$$

The 1-dimensional manifold  $T$  is the cosmic time, the 3-dimensional one  $S$  is the cosmic space. By eq.(2.1) the tangent space at a point  $p \in M$  is

$$M_p = T_p \oplus S_p. \quad (2.2)$$

In accordance with this, we have for a tensor  $K = K^{\mu\nu}$  of rank 2 the following decomposition:

$$K = K_T + K_S + K_{TS} + K_{ST}, \quad (2.3)$$

where

$$K_T = K_{TT} = K^{00}, \quad K_S = K_{SS} = K^{ij}, \quad K_{TS} = K^{0i}, \quad K_{ST} = K^{i0}. \quad (2.4)$$

For the metric tensor we have

$$g = g_T + g_S, \quad g_{TS} = g_{ST} = 0, \quad (2.5)$$

which implies  $T \perp S$ . Furthermore,

$$g_T = dt \otimes dt, \quad g = dt \otimes dt - \tilde{g}_t, \quad (2.6)$$

or

$$g = dt^2 - \tilde{g}_{tij} dx^i dx^j. \quad (2.7)$$

### 3 The state of the universe

The state of the universe is

$$\omega = (\omega_{st}, \omega_m); \quad (3.1)$$

the state of the spacetime is

$$\omega_{st} = (\tilde{g}, \tilde{g}'), \quad (3.2)$$

where prime denotes the derivative with respect to the cosmic time  $t$ ; the state of the matter is

$$\omega_m = (\Psi, \cdot\Psi), \quad (3.3)$$

where  $\Psi$  is a state vector. Thus

$$\omega = (\tilde{g}, \tilde{g}', (\Psi, \cdot\Psi)). \quad (3.4)$$

### 4 Equations of motion

The equations of motion are

$$G_S = \omega_m(T_S) = (\Psi, T_S\Psi), \quad \text{or} \quad G^{ij} = \omega_m(T^{ij}), \quad (4.1)$$

$$H_t\Psi_t = \varepsilon_t\Psi_t, \quad (4.2)$$

where  $G$  is the Einstein tensor,  $T$  is the Schrödinger operator of the energy-momentum tensor,  $H_t$  is the Schrödinger Hamiltonian:

$$H_t = \int_S \mu(ds) T_{Ts,t}(\partial/\partial t, \partial/\partial t), \quad T_T(\partial/\partial t, \partial/\partial t) = T^{00}. \quad (4.3)$$

Eq. (4.1) is the restricted Einstein equation. The equations for  $T, TS$ , and  $ST$ , i.e., for  $0\mu, \mu 0$  components are not used: The components  $G^{0\mu} = G^{\mu 0}$  do not involve  $\tilde{g}''$ , so that the corresponding equations are violated at quantum jumps, i.e., jumps of  $\omega_m$ . This violation is a manifestation of quantum holism connected with the character of quantum state.

To ensure against misunderstanding, it should be stressed that the restriction of the Einstein equation does not violate the covariance (or, more precisely, the geometric character) of the theory developed — in view of the structure of the spacetime, eq. (2.1).

Eq. (4.2) realizes the cosmic energy determinacy principle.

There are  $6+1=7$  equations (4.1),(4.2) for  $6+1=7$  quantities  $\tilde{g}_{ij}, \Psi$ .

## 5 The equation for state vector between jumps

Let us consider eq.(4.2) in more detail. We have for the Hamiltonian

$$H_t = \sum_l \varepsilon_{lt} P_{lt}, \quad (5.1)$$

$P_{lt}$  being the projector for a level  $l$ . Then eq.(4.2) boils down to

$$P_{lt} \Psi_t = \Psi_t. \quad (5.2)$$

Designating  $P_l = P$ , we have between jumps

$$\Psi = P\Psi, \quad \Psi + d\Psi = (P + dP)(\Psi + d\Psi), \quad (5.3)$$

whence

$$(I - P)d\Psi = (dP)\Psi. \quad (5.4)$$

This leads to

$$P(dP)\Psi = 0. \quad (5.5)$$

From the condition  $\|\Psi + d\Psi\| = 1$  it follows

$$Re\{(\Psi, Pd\Psi)\} = 0. \quad (5.6)$$

To determine  $Pd\Psi$  we put

$$Pd\Psi = \alpha(dP)\Psi \quad (5.7)$$

so that  $dP = 0$  would imply  $d\Psi = 0$ . By eqs.(5.7),(5.5)

$$Pd\Psi = 0. \quad (5.8)$$

Thus

$$d\Psi = (dP)\Psi, \quad \frac{d\Psi}{dt} = \frac{dP}{dt}\Psi. \quad (5.9)$$

The solution to eq.(5.9) is

$$\Psi_{t_2} = U(t_2, t_1)\Psi_{t_1}, \quad U(t_2, t_1) = T e^{\int_{t_1}^{t_2} dP_t}, \quad (5.10)$$

relations

$$U^+(t_2, t_1) = U^{-1}(t_2, t_1) = U(t_1, t_2) = T^+ e^{\int_{t_1}^{t_2} dP_t}. \quad (5.11)$$

being fulfilled.

## 6 Watched-pot dynamics

It is interesting to note that the result (5.9) may also be obtained in the language of quantum jumps. We have an orthogonal decomposition for  $\Psi = P\Psi$ :

$$\Psi = (P + dP)\Psi + (I - P - dP)\Psi = (P + dP)\Psi + (-dP)\Psi. \quad (6.1)$$

For the corresponding probabilities in the first order, relations

$$\|(-dP)\Psi\|^2 = 0, \quad \|(P + dP)\Psi\|^2 = 1 \quad (6.2)$$

hold, so that

$$\Psi \rightarrow (P + dP)\Psi, \quad (6.3)$$

whence eq.(5.9) results.

Such a dynamics is analogous to the watched-pot one, the role of a watching cook being played by the principle of cosmic energy determinacy.

## 7 Absence of degeneracy

For a general metric, the probability for a level  $l$  to be degenerate is zero, so that degeneracy occurs only at level crossings. Therefore we have

$$Tr P_l = 1, \quad \omega_m = (\Psi_l, \cdot \Psi_l) = Tr\{P_l \cdot \cdot\}, \quad (7.1)$$

and  $\omega_m$  reduces to  $l$ . Thus a state of the universe is

$$\omega = (\tilde{g}, \tilde{g}', l). \quad (7.2)$$

## 8 Quantum jump

Let us consider two crossing levels,  $l = 1, 2$ . The part of the Hamiltonian for these is

$$\tilde{H}_t = \varepsilon_{1t}P_{1t} + \varepsilon_{2t}P_{2t}. \quad (8.1)$$

Put crossing time  $t_{cross} = 0$ , then

$$\varepsilon_{10} = \varepsilon_{20} = \varepsilon_0. \quad (8.2)$$

We write  $\tilde{H}_t$  in the form of

$$\tilde{H} = \varepsilon_2 P + \nu P_1, \quad (8.3)$$

where

$$P = P_1 + P_2, \quad \nu = \varepsilon_1 - \varepsilon_2. \quad (8.4)$$

From eq.(8.3) it follows

$$\tilde{H}' = \varepsilon_2' P + \varepsilon_2 P' + \nu' P_1 + \nu P_1'. \quad (8.5)$$

Let us introduce the designation

$$\Delta f = f_{+0} - f_{-0}. \quad (8.6)$$

We put

$$f_0 = f_{-0}, \quad (8.7)$$

so that

$$\Delta f = f_{+0} - f_0. \quad (8.8)$$

Relations

$$\Delta \tilde{g} = 0, \quad \Delta g' = 0 \quad (8.9)$$

hold; as

$$H = H[\tilde{g}, \tilde{g}'], \quad (8.10)$$

we have

$$\Delta H = 0. \quad (8.11)$$

Indeed,

$$\Delta \tilde{H} = \varepsilon_{2+0} P_{+0} - \varepsilon_0 P_0 + \nu_{+0} P_{1+0} - \nu_0 P_{10}, \quad (8.12)$$

and

$$\varepsilon_{2+0} = \varepsilon_0, \quad P_{+0} = P_0, \quad \nu_{+0} = \nu_0 = 0, \quad (8.13)$$

which leads to eq.(8.11).

As

$$H' = H'[\tilde{g}, \tilde{g}', \tilde{g}'], \quad (8.14)$$

if  $\Delta \omega_m = 0$  then  $\Delta \tilde{g}'' = 0$  and

$$\Delta H' = 0 \quad (8.15)$$

must hold. We find

$$\Delta H' = \Delta(\varepsilon'_2 P) + \Delta(\varepsilon_2 P') + \Delta(\nu' P_1) + \Delta(\nu P'_1), \quad (8.16)$$

$$\Delta(\varepsilon_2 P') = 0, \quad \Delta(\varepsilon'_2 P) = (\Delta \varepsilon'_2) P_0. \quad (8.17)$$

As  $\lim_{t \rightarrow 0} \nu_t = 0$ , we put  $\Delta(\nu P'_1) = 0$ . Then

$$\Delta \tilde{H}' = (\Delta \varepsilon'_2) P_0 + \nu'_{+0} P_{1+0} - \nu'_0 P_{10} = 0. \quad (8.18)$$

Hence we obtain

$$\begin{aligned} (\Delta \varepsilon'_2 - \nu'_0) P_{10} &= -\nu'_{+0} P_{1+0} P_{10}, \\ (\Delta \varepsilon'_2) P_{20} &= -\nu'_{+0} P_{1+0} P_{20}. \end{aligned} \quad (8.19)$$

There are three solutions to eqs.(8.19):

$$\begin{aligned} Ia. \quad & P_{1+0} P_{20} = 0, \quad P_{1+0} P_{10} = P_{10}, \quad P_{1+0} = P_{10}, \quad P_{2+0} = P_{20}, \\ & \Delta \varepsilon'_2 = 0, \quad \nu'_{+0} = \nu'_0, \quad \varepsilon'_{2+0} = \varepsilon'_{20}, \quad \varepsilon'_{1+0} = \varepsilon'_{10}. \end{aligned} \quad (8.20)$$

$$\begin{aligned} Ib. \quad & P_{1+0} P_{20} = P_{20}, \quad P_{1+0} P_{10} = 0, \quad P_{1+0} = P_{20}, \quad P_{2+0} = P_{10}, \\ & \Delta \varepsilon'_2 = -\nu'_{+0}, \quad \Delta \varepsilon'_2 = \nu'_0, \quad \varepsilon'_{2+0} = \varepsilon'_{10}, \quad \varepsilon'_{1+0} = \varepsilon'_{20}. \end{aligned} \quad (8.21)$$

$$\begin{aligned} II. \quad & P_{1+0} P_{20} \neq 0, \quad P_{20}, \quad P_{1+0} P_{10} \neq 0, \quad P_{10}, \quad \{P_{1+0}, P_{2+0}\} \neq \{P_{10}, P_{20}\}, \\ & \Delta \varepsilon'_2 = 0, \quad \Delta \varepsilon'_2 - \nu'_0 = 0, \quad \nu'_{+0} = 0, \quad \varepsilon'_{2+0} = \varepsilon'_{1+0} = \varepsilon'_{20} = \varepsilon'_{10}. \end{aligned} \quad (8.22)$$

The solutions *Ia* and *Ib* in fact coincide and describe a simple crossing, i.e., that without tangency. In that case there is no jump.

The solution *II* describes a crossing with tangency. In case of such a crossing, the probability for  $\{P_{1+0}, P_{2+0}\} = \{P_{10}, P_{20}\}$  is equal to 0, so that if there is a tangency, a jump occurs with the probability 1.

Thus a criterion for the quantum jump is the level tangency.

We have

$$\begin{aligned}\omega_{t_{tang}} &= (\tilde{g}_{t_{tang}}, \tilde{g}'_{t_{tang}}, l_{t_{tang}}) \rightarrow \omega_{t_{tang}+0} = (\tilde{g}_{t_{tang}}, \tilde{g}'_{t_{tang}}, l^j_{t_{tang}+0}), \\ W_{lj} &= Tr\{P_{l^j_{t_{tang}+0}} P_{l_{t_{tang}}}\}, \quad l = 1, 2, \quad l^j = 1, 2,\end{aligned}\tag{8.23}$$

where  $W_{lj}$  is the probability for the jump  $l \rightarrow l^j$ .

## 9 Indeterministic dynamics

The indeterministic dynamics constructed above is very simple in the conceptual sense (but only in that sense!): Its only ingredients are the state of the universe

$$\omega = (\tilde{g}, \tilde{g}', l),\tag{9.1}$$

the restricted Einstein equation

$$G_S = Tr\{P_l T_S\}\tag{9.2}$$

for time evolution between jumps, and jumps at the level tangency

$$(\tilde{g}, \tilde{g}', l) \rightarrow (\tilde{g}, \tilde{g}', l^j), \quad W_{lj} = Tr\{P_{l^j} P_l\}.\tag{9.3}$$

The dynamics is unified: Both deterministic part and quantum jumps follow from the principle of cosmic energy determinacy. Further still, strange though it might seem, all the dynamics may be described in terms of quantum jumps.

Indeterministic dynamics involves some subtleties, therefore there is good reason to describe it in more detail.

For  $t_1, t_2$  between two jumps,  $t_2 > t_1$ , we introduce a designation for deterministic time evolution:

$$\omega_{t_2} = D(t_2, t_1)\omega_{t_1}.\tag{9.4}$$

In addition, we introduce an auxiliary deterministic time evolution with a fixed initial state of the matter:

$$\omega_{t_2 t_1} = D_{t_1}(t_2, t_1)\omega_{t_1}, \quad \omega_{mt} = \omega_{mt_1} \quad for \quad t_1 \leq t \leq t_2.\tag{9.5}$$

For  $t = t_{cross} = 0$  we have

$$\omega_0 = \{\tilde{g}_0, \tilde{g}'_0, P_{l_0}\}, \quad l = 1 \text{ or } 2.\tag{9.6}$$

Then for  $\delta t > 0$

$$\omega_{\delta t 0} = D_0(\delta t, 0)\omega_0 = \{\tilde{g}_{\delta t 0}, \tilde{g}'_{\delta t 0}, P_{l_0}\}.\tag{9.7}$$

For this state, the Hamiltonian  $\tilde{H}$  is

$$\tilde{H}_{\delta t 0} = \varepsilon_{1 \delta t 0} P_{1 \delta t 0} + \varepsilon_{2 \delta t 0} P_{2 \delta t 0},\tag{9.8}$$

and we put for  $t = \delta t + 0$

$$\omega_{\delta t+0} = \{\tilde{g}_{\delta t 0}, \tilde{g}'_{\delta t 0}, P_{l^j \delta t 0}\}, \quad l^j = 1 \text{ or } 2. \quad (9.9)$$

For  $t = +0$ , a true state is by definition

$$\omega_{+0} = \omega_{+00} = \lim_{\delta t \rightarrow +0} \omega_{\delta t+0} = \{\tilde{g}_0, \tilde{g}'_0, P_{l^j +0}\}, \quad P_{l^j +0} = \lim_{\delta t \rightarrow +0} P_{l^j \delta t 0}. \quad (9.10)$$

This gives the explicit definition for  $P_{l^j +0}$ .

For  $t > 0$ , the true state is given by

$$\omega_t = \lim_{\delta t \rightarrow +0} D(t, \delta t) \omega_{\delta t+0}. \quad (9.11)$$

It is important to stress that

$$\omega_t \neq D(t, +0) \omega_{+0}. \quad (9.12)$$

The value

$$\omega_{+0} = \lim_{t \rightarrow +0} \omega_t \quad (9.13)$$

coincides with that given by eq.(9.10).

## 10 Deterministic approximation

The relations

$$div G = G^{ij}_{;i} = 0, \quad (10.1)$$

$$div T_H = 0 \quad (10.2)$$

hold ( $H$  denotes the Heisenberg picture).

In the deterministic approximation, i.e., with no quantum jumps we have

$$\Psi_H = const. \quad (10.3)$$

Then

$$div(\Psi, T\Psi) = div(\Psi_H, T_H\Psi_H) = (\Psi_H, div T_H\Psi_H) = 0, \quad (10.4)$$

and the (unrestricted) Einstein equation

$$G = (\Psi, T\Psi) \quad (10.5)$$

may be fulfilled with

$$G^{0\mu} = (\Psi, T^{0\mu}\Psi) \quad (10.6)$$

as initial conditions.

Thus in the deterministic approximation, quantum holism does not manifest itself.



## References

- [1] Vladimir S. Mashkevich, *Indeterministic Quantum Gravity* (gr-qc/9409010, 1994).
- [2] R.P. Feynman, *Lecture on gravitation* (California Institute of Technology, 1962/63).
- [3] Roger Penrose, *Gravity and state vector reduction, in: Quantum Concepts in Space and Time*, ed. C. J. Isham and R. Penrose (Oxford, Oxford University Press, 1986);  
*The Emperor's New Mind* (Oxford University Press, 1989).
- [4] F. Károlyházy, A. Frenkel, and B. Lukács, *On the Possibility of Observing the Eventual Breakdown of the Superposition Principle, in: Physics as natural philosophy, essays in honor of László Tisza on his seventy-fifth birthday*, ed. A. Shimony and H. Feshbach (MIT Press, Cambridge, USA, 1982);  
*On the possible role of gravity in the reduction of the wave function, in: Quantum Concepts in Space and Time*, ed. C. J. Isham and R. Penrose (Oxford, Oxford University Press, 1986).